

Fractality in Persistence Decay and Domain Growth during Ferromagnetic Ordering: Dependence upon initial correlation

Saikat Chakraborty and Subir K. Das*

Theoretical Sciences Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Jakkur P.O, Bangalore 560064, India

(Dated: May 3, 2016)

Dynamics of ordering in Ising model, following quench to zero temperature, have been studied via Glauber spin-flip Monte Carlo simulations in space dimensions $d = 2$ and 3 . One of the primary objectives has been to understand phenomena associated with the persistent spins, viz., time decay in the number of unaffected spins, growth of the corresponding pattern and its fractal dimensionality, for varying correlation length in the initial configurations, prepared at different temperatures, at and above the critical value. It is observed that the fractal dimensionality and the exponent describing the power-law decay of persistence probability are strongly dependent upon the relative values of nonequilibrium domain size and the initial equilibrium correlation length. Via appropriate scaling analyses, these quantities have been estimated for quenches from infinite and critical temperatures. The above mentioned dependence is observed to be less pronounced in higher dimension. In addition to these findings for the local persistence, we present results for the global persistence as well. Further, important observations on the standard domain growth problem are reported. For the latter, a controversy in $d = 3$, related to the value of the exponent for the power-law growth of the average domain size with time, has been resolved.

I INTRODUCTION

Kinetics of phase transitions [1–4] remains an active area of research for several decades. In this area, typically one is interested in the nonequilibrium dynamics related to the evolution of a system to a new equilibrium state, having been quenched from a configuration prepared outside the coexistence curve to inside it, via the variation of temperature (T), pressure, etc. In this work, our focus is on the paramagnetic to ferromagnetic transition [5]. When a system is quenched, via variation of T , from the paramagnetic phase to ferromagnetic one, domains rich in like spins form and grow with time [2]. Aspects that drew attention of researchers, in this problem, are understanding of domain patterns [2], growth of domains [2], aging properties of the evolution [6–8], as well as the pattern (and corresponding dynamics) exhibited by atomic magnets (or spins) that did not change orientation till time t , referred to as persistent spins [4, 9–25]. This work deals with issues related to domain growth and persistence.

During the process of ferromagnetic ordering (where the order parameter is a nonconserved quantity), the average domain size, ℓ , increases as [2]

$$\ell \sim t^\alpha, \quad (1)$$

where α , the growth exponent, may have dependence upon system dimensionality (d) based on the order-parameter symmetry. This growth occurs via motion and annihilation of defects, facilitated by change in orientation of the spins, S_i , the subscript i being an index related to an atom or spin, typically considered to be located on a regular lattice. In this work we study the spin-1/2 Ising model, to be defined later,

for which defects are the domain boundaries. In this case, S_i is a scalar quantity which gets affected only via (complete) flipping or change in sign. For this model, the theoretical expectation for α is same in both $d = 2$ and 3 .

The persistence probability, P , defined as the fraction of unaffected spins, typically decays as [4]

$$P \sim t^{-\theta}, \quad (2)$$

where θ is expected to have dependence upon d . The persistent spins exhibit interesting fractal pattern with dimensionality [17] d_f whose dependence upon θ will be introduced later. Unless mentioned otherwise, all our results on this issue correspond to local persistence, probability for which, as already mentioned, is calculated by counting unaffected “microscopic” spins. There has also been interest in the calculation of such probability by dividing the system into blocks of linear dimension ℓ_b and counting the persistence of coarse-grained or block spin variables [14, 15]. In the limit $\ell_b \rightarrow a$, the microscopic lattice constant, such block persistence probability, P_b , will correspond to P , the “local or site persistence” probability. On the other hand, for $\ell_b \rightarrow \infty$, one obtains “global persistence” probability, further discussion and results for which will be presented later.

For Ising model, values [2, 17, 19] of α , θ and d_f are accurately estimated via Monte Carlo (MC) simulations, in $d = 2$, for quenches from initial temperature $T_i = \infty$ to the final value $T_f = 0$. It is reasonably well established [2, 4, 11–19, 24] that, in this case, the values of α , θ and d_f are $1/2$, 0.225 , and 1.58 . However, the conclusions, if exists, on the corresponding numbers for $d = 3$ are questionable [26–28]. Recent focus, on the other hand, for persistence as




FIG. 1. Log-log plots of local persistence probability, $P(t)$, vs t , for quenches (of the Glauber Ising model) from different values of initial temperature T_i ($\geq T_c$, the critical temperature), to the final value $T_f = 0$. All results correspond to space dimension $d = 2$ and square lattice, with linear dimension of the square box being $L = 2048$, in units of the lattice constant a . The lines represent various power-law decays, values of the exponents being mentioned in appropriate places.

well as for other aspects of coarsening, has been on [23–25, 29–31] quenches from temperatures providing large equilibrium correlation length ξ . In this context, in a recent work [24], we have explored the initial correlation dependence of α and θ . Our observation was, while α is insensitive to the variation of T_i (at least in $d = 2$), P (and thus θ) is strongly influenced by the choice of the latter, viz., we obtained for $d = 2$ and 3, $\theta = \theta_c \simeq 0.035$ and $\simeq 0.105$ for $T_i = T_c$, the critical temperature (see Fig. 1 for $d = 2$). The numbers quoted above are significantly different from those for $T_i = \infty$. For intermediate temperatures, as seen in Fig. 1, two step decays can be noticed. A slower decay was observed for $\ell < \xi$. The corresponding exponent θ_I approaches θ_c as $\xi \rightarrow \infty$, i.e., when $T_i \rightarrow T_c$. For $\ell \gg \xi$, behavior consistent with $T_i = \infty$ was obtained. This implies, dynamics of the spins is strongly influenced by the relative values of nonequilibrium domain length ℓ and the equilibrium correlation length ξ in the initial configuration. The overall time decay of P , for all T_i , was empirically constructed to be [24]

$$P(t)x^{2\theta} = A \left(\frac{x}{g(x) + x} \right)^\phi; \quad x = \ell/\xi, \quad (3)$$

with

$$g(x) = \frac{C_0}{1 + C_1 x^\psi}, \quad (4)$$

where A is the amplitude of the long time decay, $\phi =$

$(\theta - \theta_I)/\alpha$, $\psi \simeq 2$, whereas C_0 and C_1 are dimension dependent constants.

An extension of a study [17] (via a different model in $d = 1$) predicts

$$d_f = d - z\theta, \quad (5)$$

where z , to be more formally defined later, is a dynamical exponent related to the growth of the persistence pattern. From previous studies [24, 31], even though it has been reported that the decay of P is disconnected with the growth of ℓ , z and α may be related. Nevertheless, since such a connection is unclear, to gain knowledge about the variation of d_f , as a function of T_i , estimation of z is needed. Even if such a connection exists, as mentioned, the value of α in $d = 3$ is not unambiguous. In this dimension, the theoretically [2] expected value of α ($= 1/2$) disagrees with some computer simulations [14] which report numbers close to $1/3$. This difference can possibly [26] be due to long transient period. Thus, lengthy simulation runs with large systems are needed. It will be interesting to see if such long simulation, luck favoring, can provide the theoretically expected value. If yes, in that time regime, do we see change in other quantities as well?

In this work, our objective thus, is to estimate d_f , z , α and θ , for $T_i = \infty$ and $T_i = T_c$, in space dimensions 2 and 3, for quenches to $T_f = 0$. For the ease of reading, in TABLE 1 we provide a list of values of these quantities, obtained from computer simulations. While the ones with asterisks, to the best of our knowledge, will be calculated (or the simulation results will be shown to be consistent with those theoretical expectations) for the first time, the numbers appearing inside the parentheses are improvements over the existing ones that appear outside. We will start presenting results with the objective of calculating d_f . Other quantities will be needed for this purpose and will be estimated in due course.

TABLE 1. List of some nonequilibrium exponents for Ising model.

Case	α	z	θ	d_f
$d = 2, T_i = \infty$	$1/2$	2	0.225	1.58 (1.53)
$d = 2, T_i = T_c$	$1/2$	2*	0.035	1.92*
$d = 3, T_i = \infty$	$1/3$ (1/2)	2*	0.18 (0.15)	2.65*
$d = 3, T_i = T_c$	$1/2$	2*	0.105	2.77*

In Fig. 2 we show persistence snapshots for $T_i = \infty$ and T_c , both from $t = 10^4$ Monte Carlo steps (MCS), this time unit to be defined soon, for $d = 2$ Ising model. It is clear that the patterns are different and so, different values of d_f are expected. In Fig. 3, snapshots from an intermediate temperature $T_i = 2.4$ ($> T_c$), for $d = 2$, are presented. The first frame corresponds to a time falling in the slower decay regime of




FIG. 2. Snapshots of the persistent spins are shown for quenches from $T_i = \infty$ and T_c , to $T_f = 0$. The results correspond to $d = 2$, $L = 2048$ and $t = 10^4$ MCS. In both the cases only parts of the boxes are shown where the persistent spins are marked in black.




FIG. 3. Snapshots of the persistent spins from different times, mentioned on the figure, are shown for $T_i = 2.4$ and $T_f = 0$. Other details are same as Fig. 2.

Fig. 1 (for the corresponding temperature), whereas the second one is from the faster decay regime, implying $\ell \gg \xi$. The earlier time snapshot resembles the $T_i = T_c$ picture of Fig. 2 and the second one has similarity with $T_i = \infty$ pattern. This justifies our focus only on these two limiting initial temperatures with $\xi = 0$ and ∞ , rather than exploring a wide temperature range, to accurately quantify d_f and z .

The rest of the paper is organized as follows. In the next section we describe the model and method. Section III provides a brief overview of an earlier work. Results are presented in section IV. Finally, section V concludes the paper with a brief summary and outlook.

II MODEL AND METHOD

As already mentioned, we study the Ising model [5], on square or simple cubic lattice systems, depending upon the dimensionality, with nearest neighbor interactions. The Hamiltonian for the model is given by

$$H = -J \sum_{\langle ij \rangle} S_i S_j; \quad S_i = \pm 1, \quad (6)$$

where J is the interaction strength (> 0) and $\langle ij \rangle$ implies interaction among nearest neighbors. The values of T_c for this model in $d = 2$ and 3 are respectively [32] $\simeq 2.27J/k_B$ and $\simeq 4.51J/k_B$, k_B being the Boltzmann constant.

Kinetics in this model was introduced via Glauber spin-flip mechanism [32, 33]. In this MC approach, a trial move consists of changing the sign of a randomly chosen spin. Since our quenches were done to $T_f = 0$, a move was accepted only if it had reduced the energy. Needless to say, initial configurations were prepared at nonzero T values. In that case, the Metropolis criterion for the acceptance of a move was implemented via appropriate calculation of the Boltzmann factor [32] and its comparison with a random number, ranging between 0 and 1, whenever the move brought an increment in the energy. For preparation of initial configurations at temperatures very close to T_c , in addition to the Glauber mechanism, we have applied Wolff algorithm [34] as well, which facilitates faster equilibration. Time, in our simulations are measured in units of MCS, each MCS consisting of L^d steps, L being the linear dimension of a square or cubic box. Periodic boundary conditions were applied in all directions. Final results are presented after averaging over multiple initial realizations, the number ranging from 20 to 70. In $d = 2$ all results are for $L = 2048$. In $d = 3$, the results for $T_i = \infty$ are for $L = 512$ and for $T_i = T_c$, we presented results from $L = 400$ and 256 .

III AN OVERVIEW OF THE BACKGROUND ON FRACTALITY OF PERSISTENCE PATTERN

In this section we provide a discussion on the theoretical background for fractality of the structures formed by persistent spins, following the work by Manoj and Ray [17].

From a density correlation function, $D(r, t)$, isotropic in an unbiased system, total mass or number of particles in a circular or spherical (depending upon dimensionality) region of radius R can be obtained as

$$M(R, t) \sim \int_0^R D(r, t) r^{d-1} dr, \quad (7)$$

r ($= |\vec{r}|$) being the scalar distance of a point in that region from the central one. An appropriate correlation function in the present context is

$$D(r, t) = \frac{\langle \rho(\vec{r}_0, t) \rho(\vec{r}_0 + \vec{r}, t) \rangle}{\langle \rho(\vec{r}_0, t) \rangle}, \quad (8)$$

with ρ being unity at a space point if the spin there did not flip till time t and zero otherwise. The average

order parameter for the persistent pattern is

$$\langle \rho(\vec{r}, t) \rangle = \frac{\int d\vec{r} \rho(\vec{r}, t)}{\int d\vec{r}} = P(t). \quad (9)$$

This being a nonconserved (time dependent) quantity and, since, in the definition of $D(r, t)$, the average value is not subtracted from ρ , decorrelation here means, decay of $D(r, t)$ to a “non-zero” value ($=P(t)$), for $t < \infty$. The distance, $\ell_p(t)$, at which $D(r, t)$ reaches this plateau is the characteristic length scale of the pattern. In that case, there may exist scaling of the form

$$\frac{D(r, t)}{P(t)} \equiv f(r/\ell_p). \quad (10)$$

For $x (\equiv r/\ell_p) > 1$, f should be unity. On the other hand, for fractal dimension d_f and $x < 1$, one should have

$$f(x) \sim x^{d_f - d}, \quad (11)$$

since

$$M \sim x^{d_f}. \quad (12)$$

Considering that $P(t)$, the plateau value, decays in a power-law fashion, a power-law behavior of $f(x)$ is indeed expected, once scaling is achieved. A continuity, at $r = \ell_p$, in such a situation demands

$$t^{(d_f - d)/z} = t^{-\theta}, \quad (13)$$

providing Eq. (5), where z is the dynamic exponent characterizing the growth of the persistence pattern, mentioned before, as

$$\ell_p \sim t^{1/z}. \quad (14)$$

For this model, as mentioned, value of α has been estimated [24] for various T_i values in $d = 2$. However, a priori it is unclear whether there is a general validity of the relation

$$z\alpha = 1. \quad (15)$$

Then it is necessary to calculate both z and θ , for correlated and uncorrelated initial configurations, to validate Eq. (5). On the other hand, as already mentioned, the value of α is ambiguous in $d = 3$.

IV RESULTS

In Fig. 4 we show $D(r, t)$ as a function of r for $T_i = 2.4$, from two different times, mentioned on the figure, for $d = 2$. As expected, the correlation function decays to different constant value, $P(t)$, at different length ℓ_p , for different times. Before decaying to the




FIG. 4. Density correlation functions, $D(r, t)$, related to the persistent spins, are plotted vs r . Results are presented from two different times, for $T_i = 2.4$ and $T_f = 0$. The system dimensionality is $d = 2$ and value of L is 2048.

plateau, the early time data appear to obey a power-law. The later time data, for smaller r , follows the same power-law before crossing over to another, faster, power-law decay. This implies, there exist two length scales in the problem, below and beyond the equilibrium scale ξ . Inside the larger structure, the small length scale structure remains hidden, which will become irrelevant in the long time limit. For $\xi = \infty$, i.e., $T_i = T_c$, however, the latter will be the only structure and remain for ever. The exponent for large r , for $T_c < T_i < \infty$ and $t \gg 0$, should be related to the d_f value for $T_i = \infty$ case whereas, in case of small r , the exponent should be connected to d_f for $T_i = T_c$ case. Below we focus on these two cases, i.e., $T_i = \infty$ and $T_i = T_c$, separately, first for $d = 2$, followed by $d = 3$. As need occurs, we will present results related to α , θ and z .

In the main frame of Fig. 5 we present a scaling exercise [17, 19] for $D(r)$ where we have plotted $f(x)$ as a function of x , using data from different times after quench, for $T_i = T_c$ and $d = 2$. Scaling appears good and gets better with the progress of time. On this log-log plot, look of the data appear, before decaying to unity, linear, implying a power-law decay. The exponent appears to be $\simeq 0.09$. In the inset of this figure, we show analogous exercise for $T_i = \infty$. Even though this case in this dimension was studied by Jain and Flynn [19], for the sake of comparison and completeness, we present it here from our own simulations. In this case, the exponent for the power-law decay appears consistent with 0.45. Then, in $d = 2$, for $T_i = \infty$, the fractal dimensionality is 1.55 and for $T_i = T_c$, the number is 1.91, if Eq. (11) is valid.

In Fig. 6 we show the plots of ℓ_p vs t in $d = 2$, for (a) $T_i = \infty$ and (b) $T_i = T_c$, on log-log scales. In




FIG. 5. Scaling analysis of $D(r,t)$ for the $d = 2$ Ising model, with $T_i = T_c$ and $T_f = 0$, where $f(x)$ is plotted vs $x = r/\ell_p$, using data from different times after the quench, on log-log scale. The solid line corresponds to a power-law decay with an exponent 0.09. Inset: Same as the main frame but for $T_i = \infty$. The solid line here has the power-law decay exponent 0.45. The value of L is 2048 for all the results.

both the cases the data appear consistent with $z = 2$, validating Eq. (15) (note that α is established to be $1/2$ in $d = 2$). Nevertheless, we intend to make more accurate quantification. For this purpose, in the insets of these figures we have shown instantaneous exponents (dash-dotted lines), z_i , calculated as [35]

$$\frac{1}{z_i} = \frac{d \ln \ell_p}{d \ln t}, \quad (16)$$

vs $1/\ell_p$. In both the cases we obtain the value of z via linear extrapolation (see the consistency of the simulation data with the solid line) to $\ell_p = \infty$. For $T_i = \infty$, from this exercise, we quantify $z = 2.15$ and for $T_i = T_c$, we obtain the number $z = 2.02$ (see the dashed horizontal lines). These numbers, in addition to verifying Eq. (15), are also consistent with the numbers obtained via least square fitting of the ℓ_p vs t data to the form

$$\ell_p = \ell_z^0 + A_z t^{1/z}, \quad (17)$$

where ℓ_z^0 and A_z are positive constants. This consistency may imply, early time corrections to the exponents are insignificant. Note here that, in absence of any correction, one expects [36, 37]

$$\frac{1}{z_i} = \frac{1}{z} \left[1 - \frac{\ell_z^0}{\ell_p} \right], \quad (18)$$

a linear behavior of $1/z_i$, when plotted vs $1/\ell_p$, with slope $-\ell_z^0/z$. A positive slope in both the insets is




FIG. 6. (a) Log-log plot of persistence length scale, ℓ_p , as a function of t , for $d = 2$ Ising model, following quench from $T_i = \infty$ to $T_f = 0$, with $L = 2048$. Inset: Instantaneous exponent, z_i , obtained using the data in main frame, is plotted vs. $1/\ell_p$. (b) Same as (a), but here $T_i = T_c$. The solid lines in the main frame of both (a) and (b) correspond to power-law growths with exponent $1/2$. In the insets, the horizontal dashed lines correspond to our estimates for z , whereas solid lines are guides to the eyes.

due to the fact that we have presented inverse of the quantity discussed in Eq. (18). Using these values of z , and numbers for θ , mentioned earlier, in Eq. (5), we obtain $d_f = 1.93$ for $T_i = T_c$ and $d_f = 1.51$ for $T_i = \infty$. These values, within computational errors, are consistent with the conclusions from Fig. 5. Next we present results from $d = 3$.

In $d = 3$, we start by presenting results for the growth of ℓ , considering the controversy [14, 24, 26] on the value of α discussed above. In $d = 2$, we avoided

presenting results on this aspect with the understanding that the issue there is well settled. Nevertheless, in the context of global persistent decay, we will make indirect conclusion about it. Here note that the estimation of ℓ was done from the first moment of domain size distribution, $p(\ell_d, t)$, as

$$\ell(t) = \int \ell_d p(\ell_d, t) d\ell_d, \quad (19)$$

where ℓ_d is the distance between two domain boundaries in a particular direction. The main frame of Fig. 7(a) shows a plot of ℓ vs t , on log-log scale, for quenches of the $d = 3$ Ising system from $T_i = \infty$ to $T_f = 0$. There exists an intermediate time regime, extending over more than two decades, during which the simulation data show consistency with an exponent $\alpha = 1/3$, in agreement with previous results [14]. However, as discussed and a trend demonstrated in Ref. [26], the discrepancy in the earlier reports from the theoretical number $1/2$ can be due to long transient. Thus, long simulation runs with large systems are necessary. We have simulated a system with $L = 512$ for time longer than any of the previous works, to the best of our knowledge. Indeed, it appears that the long time behavior, over the longest time decade in the presented time range, is consistent with $\alpha = 1/2$. In the inset of this figure we show the instantaneous exponent

$$\alpha_i = \frac{d \ln \ell}{d \ln t}, \quad (20)$$

as a function of $1/\ell$. This provides an accurate picture, the long time exponent being within 5% of the theoretical value. One may then ask, is the value of θ going to change, beyond this crossover time? Even if θ has no dependence on the value of α , such a change may still occur. Note that conclusion on the value of θ , in earlier works [16, 24], were drawn from runs shorter than this. Indeed, a jump in θ_i , calculated from

$$\theta_i = -\frac{d \ln P}{d \ln t}, \quad (21)$$

occurs (see Fig. 7(b) and the corresponding inset) from an early time value of $\simeq 0.18$ to $\simeq 0.15$. This may, of course, be due to statistical or other reasons. However, since the jump in α_i occurs around the same time as the one for θ_i and fluctuation is seen around stable mean values, in both α_i and θ_i , we accept this as the correct number for θ in the asymptotic time limit.

Whether due to lattice anisotropy [14] or anything else, the solution to overcome such long transient is certainly related to being able to access large length scales. For T_i close to T_c , since this is automatically the case, due to large initial correlation, we expect an

enhanced value of α from early time. Corresponding ℓ vs t data are presented in Fig. 8. On the log-log scale, this data set shows consistency with $\alpha = 0.45$. Here we mention that study for $T_i = T_c$ has additional problems related to longer equilibration time at the initial temperature and stronger finite-size effects during the nonequilibrium evolution [24]. The latter remark can be appreciated from the plot in Fig. 8 where a bending of the data set (from the power-law behavior) is visible from $t = 10^3$. This should be compared with the corresponding data in Fig. 7(a) for $T_i = \infty$. Thus, accessing very large length scales, without finite-size effects, for $T_i = T_c$, is extremely difficult. The P vs t data, shown in the inset of Fig. 8, exhibit consistency [24] with $\theta_c \simeq 0.105$. Since, α is very close to $1/2$ already, we do not expect much change in θ_c even in true asymptotic length or time limit. The calculations of α_i and θ_i in this case provide numbers consistent with the ones quoted above.

Next, we come back to the issue of fractality. For $d = 3$ Ising model, our results in this context are entirely new irrespective of the value of T_i .

Fig. 9(a) is analogous to the inset of Fig. 5 ($T_i = \infty$ results for $f(x)$) but for $d = 3$. The corresponding $f(x)$ vs x scaling plot for $T_i = T_c$ in $d = 3$ is presented in Fig. 9(b). Again, for both $T_i = \infty$ and $T_i = T_c$, good data collapse are obtained for results from different times, in these scaling plots. For both values of T_i , we have used data sets lying in the time ranges that provide consistency with the expected theoretical number for α . In the relevant region, the $T_i = T_c$ results have power-law decay with exponent 0.24. In case of $T_i = \infty$, the value of this exponent is approximately 0.38. These numbers imply $d_f = 2.76$ and 2.62 for $T_i = T_c$ and $T_i = \infty$, respectively.

In Fig. 10 we show ℓ_p vs t plots from $d = 3$ for (a) $T_i = \infty$ and (b) $T_i = T_c$. In the long time limit, the results, in both (a) and (b), appear consistent with growth having $z = 2$. This is in agreement with Eq. (15). From the log-log plot for $T_i = \infty$, like ℓ vs t , a long time transient is clearly visible. To quantify z more accurately (in the $t \rightarrow \infty$ limit), for both $T_i = \infty$ and $T_i = T_c$, we have shown the instantaneous exponents, vs $1/\ell_p$, in the insets. From there, we extract $z = 2.1$ for $T_i = \infty$ and 2.15 for $T_i = T_c$. Along with the above mentioned numbers for z , using the values of θ for quenches from $T_i = T_c$ and $T_i = \infty$, we obtain $d_f \simeq 2.78$ and 2.68 . These numbers are consistent with those obtained from the scaling plots in Fig. 9, providing higher confidence on our estimation of θ from long time limit, for $T_i = \infty$. An interesting exercise here would have been to plot $z_i \alpha_i$ vs t . However, a constant value of unity cannot be obtained because of the fact that ℓ and ℓ_p have different initial off-sets. This is evident from the pictures in the insets of Fig. 7(a) and Fig. 10(a). While for the time dependence of ℓ , a long transient with $\alpha \simeq 1/3$ is visible, this is




FIG. 7. (a) Log-log plot of ℓ vs t , in $d = 3$, for $T_i = \infty$. The solid lines correspond to different power laws, exponents for which are mentioned. The inset shows instantaneous exponent α_i , as a function of $1/\ell$. The dashed horizontal lines represent exponent values 0.36 and 0.48. (b) Log-log plot of $P(t)$ vs t , for $d = 3$ and $T_i = \infty$. The inset shows instantaneous exponent θ_i vs $1/\ell$. Horizontal dashed lines are for $\theta = 0.176$ and 0.15.

not so for the time dependence of ℓ_p . Thus, because of the off-set related reason mentioned above, $z_i \alpha_i = 1$ is expected to be valid only in the $t \rightarrow \infty$ limit.

Finally, we turn our attention to the block persistence which was introduced by Cueille and Sire [14]. The corresponding probability P_b , as already mentioned, is related to the change in the order-parameter variable obtained by coarse-graining the site or microscopic spin variables over a block of linear size ℓ_b . It is expected that the decay of this probability will be significantly slower than the site or local persistence




FIG. 8. Log-log plot of ℓ vs t , for $d = 3$, $L = 400$ and $T_i = T_c$. The solid line corresponds to a power-law growth with exponent 0.45. The inset shows a plot of P vs t , on log-log scale. The solid line there represents a power-law decay with exponent 0.105.

probability, to which the former should cross over only for $\ell > \ell_b$. This two time-scale behavior is desirable by considering that, in the early time regime, a slower decay is forced by the fact that a sign change in block spin variable happens only when ℓ becomes comparable to ℓ_b and in the large ℓ limit, the blocks effectively appear as sites. It is expected then that a scaling should be obtained as [14]

$$P_b \ell_b^{\theta_0/\alpha} \equiv h(t/\ell_b^{1/\alpha}), \quad (22)$$

where θ_0 is the exponent of the early part of the decay or global persistence exponent in the sense that when $\ell_b \rightarrow \infty$, this is the only exponent. In $d = 2$, we will see that the best scalings, in accordance with Eq. (22), are obtained for $\alpha = 1/2$, irrespective of the value of T_i . In $d = 3$, on the other hand, due to long transient in the dynamics, we avoid presenting these results.

In addition to the above mentioned understandings, calculation of persistence probability via such blocking may have advantage for quenches to nonzero temperature. Note that for $T_f \neq 0$, thermal fluctuation from bulk of the domains affects the calculation when done via standard method. Considering that domain growth occurs essentially due to spin flips along the domain boundaries, in the calculation of P , dynamics inside the domains needs to be discarded. In a method, prescribed by Derrida [12], this is done by simulating an ordered system, alongside the coarsening one, and subtracting the common flipped spins, identifiable as the bulk flips, between the two systems, from the total, thus sticking to the effects of only the boundary motion. In the block spin method, if ℓ_b is




FIG. 9. (a) Scaling function $f(x)$ is plotted vs x , for $d = 3$, $T_i = \infty$ and $T_f = 0$, using data from few different times. The solid line has a power-law decay with exponent 0.38. (b) Same as (a) but, instead of $T_i = \infty$, we present data for $T_i = T_c$. Here the solid line has power-law decay exponent 0.24. The results were obtained for simple cubic lattice with $L = 256$.

significantly larger than ξ at T_f , thermal fluctuations will not alter the sign of block spins and in the large ℓ ($> \ell_b$) limit, as previously stated, one expects the decay to be consistent with local persistence. This saves computational time for simulating the additional systems with ordered configurations.

In Fig. 11(a) we show P_b vs t plots from $d = 2$, for a few different values of ℓ_b and $T_i = T_c$. It appears, as discussed, there exist two step decays and crossover to the faster (consistent with the local persistent decay) one is delayed with increasing ℓ_b .

In Fig. 11(b) we show a scaling exercise using the




FIG. 10. (a) Log-log plot of persistence length scale, ℓ_p , as a function of t , for $d = 3$ Ising model, following quench from $T_i = \infty$ to $T_f = 0$, with $L = 512$. Inset: Instantaneous exponent, z_i , obtained using the data in main frame, is plotted vs. $1/\ell_p$. (b) Same as (a), but here $T_i = T_c$ and $L = 256$. The solid lines in the main frame of both (a) and (b) correspond to power-law growths with exponents mentioned there. The horizontal dashed lines in the insets correspond to our estimates for z and the solid lines there are guides to the eyes.

data of Fig. 11(a) where we have plotted $h(x)$ vs $t/\ell_b^{1/\alpha}$. For obtaining collapse of data, we have adjusted θ_0 and α . The value of α used here is 0.49, that provides the best collapse. This number is certainly consistent with $1/2$, within numerical error. Early time behavior corresponds to global persistence with $\theta_0 = 0.002$ and the late time behavior is consistent with our previous estimation of $\theta_c \simeq 0.035$, for the site persistence probability. In the inset we have shown




FIG. 11. (a) Plots of block persistence probabilities, vs t , from different values of ℓ_b , for $T_i = T_c$, in $d = 2$. (b) Scaling plots of the persistence probabilities in (a). The scaling function $h(x)$ is plotted, on a log-log scale, vs $x = t/\ell_b^{1/\alpha}$. In the inset of (b) we present similar scaling plot for $T_i = \infty$. System sizes correspond to $L = 2048$. Various power-law decays are shown by solid lines with the exponent values being mentioned next to appropriate lines.

corresponding scaling results for $T_i = \infty$, for which θ_0 and θ values (mentioned on the figure) are consistent with previous findings [14]. The value of α that provides the best collapse here is 0.5. Note that in our earlier work such independence of α from T_i was directly (from the analysis of ℓ vs t data) checked for this dimension.

V CONCLUSION

We have presented results for coarsening dynamics in Ising model, with nonconserved order parameter, from space dimensions $d = 2$ and 3 . The results include domain growth law and persistence, for quenches with initial configurations of varying correlation length ξ . While presented results for persistence are mostly related to local order parameter [9–13], for the global case [14, 15] we have obtained new exponent for quench from initial temperature $T_i = T_c$, in $d = 2$. For local persistence, our results are summarized in the next paragraph.

A central objective of this paper has been to identify the differences in the patterns formed by persistent spins when systems are quenched from $T_i = \infty$ and $T_i = T_c$, to the final temperature $T_f = 0$. For both the cases, corresponding fractal dimensionalities d_f , as well as the exponent z , related to the growth of the persistent pattern, have been obtained in various dimensions. A scaling law connecting d_f , d , z and θ , predicted by Manoj and Ray [17], has been observed to be valid, irrespective of the values of d and T_i . Combining various methods, we quote, for $T_i = \infty$,

$$\begin{aligned} d_f &= 1.53 \pm 0.02, \quad d = 2, \\ d_f &= 2.65 \pm 0.03, \quad d = 3, \end{aligned} \quad (23)$$

and for $T_i = T_c$,

$$\begin{aligned} d_f &= 1.92 \pm 0.02, \quad d = 2, \\ d_f &= 2.77 \pm 0.02, \quad d = 3. \end{aligned} \quad (24)$$

On the standard domain growth problem, it is shown that the values of α in both dimensions for all initial temperatures are consistent with the theoretical expectation $\alpha = 1/2$. This number describes the growth of the persistent pattern as well, validating Eq. (15) and confirming that domain growth occurs essentially due to dynamics of spins along the domain boundaries. This resolves a controversy in $d = 3$ for which some previous computer simulations reported $\alpha = 1/3$. As mentioned in Ref. [26], this discrepancy must have been due to lack of data for extended period of time. Long simulations in our work, in addition to resolving this controversy, corrects the value of θ as well in this dimension.

In future we will address similar issues for conserved order parameter dynamics, including aging phenomena. For both conserved and nonconserved dynamics, scaling properties and form of the two-point correlation function will be an important problem for the case of correlated initial configurations.

ACKNOWLEDGEMENT

The authors thank Department of Science and Technology, Government of India, for financial support. SKD acknowledges hospitality and financial supports from International Centre for Theoretical Physics, Italy. He is also thankful to the Marie Curie Actions Plan of European Commission (FP7-PEOPLE-2013-IRSES grant No. 612707, DIONICOS).

* das@jncasr.ac.in

-
- [1] A. Onuki, *Phase Transition Dynamics*, Cambridge University Press, Cambridge, UK (2002).
 - [2] A. J. Bray, *Adv. Phys.* **51**, 481 (2002).
 - [3] R. A. L. Jones, *Soft condensed matter*, Oxford University Press, Oxford (2008).
 - [4] A. J. Bray, S. N. Majumdar and G. Schehr, *Adv. Phys.* **62**, 225 (2013).
 - [5] N. Goldenfeld, *Lecture Notes on Phase Transitions and the Renormalization Group*, Addison-Wesley, Reading, MA (1992).
 - [6] D. S. Fisher and D. A. Huse, *Phys. Rev. B* **38**, 373 (1988).
 - [7] F. Corberi, E. Lippiello and M. Zannetti, *Phys. Rev. E* **74**, 041106 (2006).
 - [8] J. Midya, S. Majumder and S. K. das, *J. Phys. : Condens. Matter* **26**, 452202 (2014).
 - [9] S. N. Majumdar, C. Sire, A. J. Bray and S. J. Cornell, *Phys. Rev. Lett.* **77**, 2867 (1996).
 - [10] S. N. Majumdar, A. J. Bray, S. J. Cornell and C. Sire, *Phys. Rev. Lett.* **77**, 3704 (1996).
 - [11] B. Derrida, V. Hakim and R. Zeitak, *Phys. Rev. Lett.* **77**, 2871 (1996).
 - [12] B. Derrida, *Phys. Rev. E* **55**, 3705 (1997).
 - [13] D. Stauffer, *Int. J. Mod. Phys. C* **8**, 361 (1997).
 - [14] S. Cueille and C. Sire, *J. Phys. A: Math and Gen.* **30**, L791 (1997).
 - [15] S. Cueille and C. Sire, *Eur. Phys. J. B* **7**, 111 (1999).
 - [16] G. Manoj and P. Ray, *Phys. Rev. E* **62**, 7755 (2000).
 - [17] G. Manoj and P. Ray, *J. Phys. A: Math and General* **33**, 5489 (2000).
 - [18] G. Manoj and P. Ray, *J. Phys. A: Math and General* **33**, L109 (2000).
 - [19] S. Jain and H. Flynn, *J. Phys. A: Math and General* **33**, 8383 (2000).
 - [20] M. Saharay and P. Sen, *Physica A* **318**, 243 (2003).
 - [21] D. Chakraborty and J. K. Bhattacharjee, *Phys. Rev. E* **76**, 031117 (2007).
 - [22] R. Paul, A. Gambassi and G. Schehr, *Europhys. Lett.* **78**, 10007 (2007).
 - [23] T. Blanchard, L. F. Cugliandolo and M. Picco, *J. Stat. Mech.* P12021 (2014).
 - [24] S. Chakraborty and S. K. Das, *European Phys. J. B* **88**, 160 (2015).
 - [25] A. Gambassi, R. Paul, G. Schehr, *J. Stat. Mech.* P12029 (2010).
 - [26] F. Corberi, E. Lippiello and M. Zannetti, *Phys. Rev. E* **78**, 011109 (2008).
 - [27] J. G. Amar and F. Family, *Bull. Am. Phys. Soc.* **34**, 491 (1989).
 - [28] J. D. Shore, M. Holzer, and J. P. Sethna, *Phys. Rev. B* **46**, 11376 (1992).
 - [29] C. Dasgupta and R. Pandit, *Phys. Rev. B* **33**, 4752 (1986).
 - [30] K. Humayun and A. J. Bray, *J. Phys. A: Math and Gen.* **24**, 1915 (1991).
 - [31] A. Sicilia, J. A. Arenzon, A. J. Bray and L. F. Cugliandolo, *Phys. Rev. E* **76**, 061116 (2007).
 - [32] D. P. Landau and K. Binder, *A Guide to Monte Carlo Simulations in Statistical Physics*, Cambridge University Press, Cambridge (2009).
 - [33] R. J. Glauber, *J. Math. Phys.* **4**, 294 (1963).
 - [34] U. Wolff, *Phys. Rev. Lett.* **62**, 361 (1989).
 - [35] D. A. Huse, *Phys. Rev. B* **34**, 7845 (1986).
 - [36] J. G. Amar, F. E. Sullivan and R. D. Mountain, *Phys. Rev. B* **37**, 196 (1988).
 - [37] S. Majumder and S. K. Das, *Phys. Rev. E* **81**, 050102 (2010).